

THE INTEGRAL REPRESENTATION FOR THE PRODUCT OF TWO PARABOLIC CYLINDER FUNCTIONS $D_\nu(x)D_\nu(-x)$ AT $\operatorname{Re} \nu < 0$ BY MEANS OF THE FUNDAMENTAL SOLUTION OF A LANDAU-TYPE OPERATOR

C. Malyshev

V.A.Steklov Institute of Mathematics at St.-Petersburg

Fontanka 27, St.-Petersburg 191011, Russia

E-mail: malyshev@pdmi.ras.ru

February 1, 2008

Abstract

The fundamental solution (Green's function) of a first order matrix ordinary differential equation arising in a Landau-type problem is calculated by two methods. The coincidence of the two representations results in the integral formula for the product of two parabolic cylinder functions $D_\nu(x)D_\nu(-x)$ at $\operatorname{Re} \nu < 0, x \in \mathbb{R}$.

PDMI Preprint 04/2001
math-ca/0106142

1 INTRODUCTION

This paper is to point out another integral representation for the product of two parabolic cylinder functions $D_\nu(x)D_\nu(-x)$ at $\text{Re } \nu < 0, x \in \mathbb{R}$. The formula to be obtained is due to equivalence of two representations for the fundamental solution (Green's function) of a first order (2×2 -matrix) ordinary differential equation related to a couple of Weber's equations. The parabolic cylinder functions have been extensively studied in classical mathematical physics [1–5] (and refs. therein). In particular, among various relations for them, the following integral representations are known for products of two parabolic cylinder functions:

$$D_\nu(z) D_{-\nu-1}(z), \quad (1.1)$$

(z real, $\text{Re } \nu < 0$, or $\text{Re } z > 0, \text{Re } \nu > -1$),

$$D_{-\nu-1}(ze^{i\frac{\pi}{4}}) D_{-\nu-1}(ze^{-i\frac{\pi}{4}}), \quad (1.2)$$

($\text{Re } z > 0, \text{Re } \nu > -1$, or $|\arg z| < \frac{\pi}{4}, \text{Re } \nu < -1$, or $-1 < \text{Re } \nu < 0$),

$$D_\nu(ze^{i\frac{\pi}{4}}) D_\nu(ze^{-i\frac{\pi}{4}}), \quad (1.3)$$

($|\arg z| < \frac{\pi}{4}, \text{Re } \nu < 0$) [1, 2, 6]. Besides, the following relations can be found in [5]:

$$-\frac{1}{z} \frac{d}{dz} (D_\nu(z) D_\nu(-z)) = \int_0^\infty J_0(zs) D_\nu^2(s) ds, \quad (2)$$

and [6]:

$$\begin{aligned} (2z)^{-1} (D_\nu(-z) D_{\nu+1}(z) - D_{\nu+1}(-z) D_\nu(z)) \\ = (2z)^{-1} \left(z - \frac{d}{dz} \right) D_\nu(z) D_\nu(-z) \\ = \int_0^\infty J_0(zs) D_\nu(s) D_{\nu+1}(s) ds. \end{aligned} \quad (3)$$

Moreover, Ref.[1] suggests to consult with [7, 8, 9] for other information concerning integral representations for the parabolic cylinder functions. Essentially, apart from (1), only the following combinations can be found in [7–16]:

$$(D_\nu(ze^{i\frac{\pi}{2}}) + D_\nu(ze^{-i\frac{\pi}{2}})) (D_{-\nu-1}(ze^{i\frac{\pi}{2}}) + D_{-\nu-1}(ze^{-i\frac{\pi}{2}})) \quad (4.1)$$

($-1 < \text{Re } \nu < 0$),

$$(D_\nu(ze^{i\frac{\pi}{2}}) - D_\nu(ze^{-i\frac{\pi}{2}})) (D_{-\nu-1}(ze^{i\frac{\pi}{2}}) - D_{-\nu-1}(ze^{-i\frac{\pi}{2}})) \quad (4.2)$$

($-2 < \text{Re } \nu < 0$),

$$D_\nu(z) (D_\nu(ze^{i\frac{\pi}{2}}) \pm D_\nu(ze^{-i\frac{\pi}{2}})) \quad (4.3)$$

($|\arg z| < \pi/4, \text{Re } \nu < 0$). There are more other representations for products of $D_\nu(z)$ by means of integrals and series in [6, 17–24]. For instance, the products at the same arguments can be found in [19, 21–23]. Certain representations through complex and contour integrals are given in [22]. Expressions for products by means of indefinite integrals are given in [19]. Recently, it has been found that the wave field of the plane wave, which is

scattered by a cone of arbitrary shape, can also be expressed for singular directions by means of the parabolic cylinder functions (the scalar and electromagnetic cases) [25].

Combining (2) and (3) we obtain the following integral for the product of two functions $D_\nu(z)$ at the opposite arguments:

$$D_\nu(z) D_\nu(-z) = -2 \int_0^\infty J_0(zs) D_\nu(s) D'_\nu(s) ds \quad (5)$$

(notice that integration of (2) agrees with (5)). In its turn, the present paper is to point out another formula for the product $D_\nu(x) D_\nu(-x)$:

$$D_\nu(x) D_\nu(-x) = 2^{-1/2} \Gamma^{-1}(-\nu) \times \int_0^\infty \exp\left(\left(\nu + \frac{1}{2}\right)t - \frac{x^2}{2} \tanh \frac{t}{2}\right) \frac{dt}{\sqrt{\sinh t}}, \quad \text{Re } \nu < 0, x \in \mathbb{R}, \quad (6)$$

which appears as a by-product of [26, 27] where a theoretical problem of condensed matter physics is studied. Specifically, the ground state average of momentum operator (so-called, mass current) in weakly inhomogeneous A-phase of superfluid helium-3 ($^3\text{He}-\text{A}$) is investigated in [26, 27]. Equation (6) appears since the Green function of the quantum-statistical model under consideration (i.e., the fundamental solution to an appropriate matrix ordinary differential equation) can be obtained in two equivalent forms: the integral [26], and the series [27]. Equating to each other, one obtains (6). The differential equation under consideration below reminds the famous physical problem of quantization for a spinning electron in a constant homogeneous magnetic field (the Landau problem) [28].

It is especially convenient to apply to (6) the Laplace method (steepest descent) in order to calculate asymptotical expansions for the mass current [27]. Therefore, Eq.(6) looks attractive since can be considered at the same footing as (1)–(5), and, hopefully, can be useful for applications, since contains only elementary functions under integration. The present paper briefly recalls the main points of [26] and [27] in Sections 2 and 3, respectively. Section 4 contains special examples, and Section 5 concludes the paper.

2 THE INTEGRAL REPRESENTATION

The master equation of our approach is given by the matrix ordinary differential equation

$$(i\omega + \mathcal{H}(x)) G(x) = e^{ix\xi} \mathbf{1}, \quad \mathcal{H}(x) \equiv \begin{pmatrix} i\frac{d}{dx} & -(x + i\Delta) \\ -(x - i\Delta) & -i\frac{d}{dx} \end{pmatrix}, \quad (7)$$

where $x \in \mathbb{R}$ is a real variable; Δ, ω, ξ are real parameters, $i \equiv \sqrt{-1}$, $\mathbf{1}$ is unit matrix, and $G(x)$ is unknown matrix. Equation (7) appears from the Dyson–Gorkov equation which describes two-point correlation functions (Green’s functions) of $^3\text{He}-\text{A}$ [26]. Under certain physically motivated conventions, Eq.(7) appears from a spatially three-dimensional general Dyson–Gorkov equation, and the variable x turns out to be due to a separation of the system into a collection of one-dimensional subsystems (spherical coordinates are chosen). The parameter Δ depends, generically, on the angle variables. The parameter ω (7) corresponds to a thermal Matsubara frequency. Solutions to the Dyson–Gorkov equation describe the physics of the model in question [26, 27].

It is convenient to place the Dirac δ -function in the R.H.S. of (7):

$$(i\omega + \mathcal{H}(x)) G(x, x') = e^{ix\xi} \delta(x - x') \mathbf{1}, \quad (8)$$

so that (7) appears after integrating (8) over x' from $-\infty$ to ∞ with $G(x) = \int G(x, x') dx'$. Let us put (8) in another equivalent form:

$$(i\frac{d}{dx} + M(x)) \tilde{G}(x, x') = e^{ix\xi} \delta(x - x') \mathbf{1}, \quad M(x) \equiv \begin{pmatrix} i\omega & x + i\Delta \\ -x + i\Delta & -i\omega \end{pmatrix}. \quad (9)$$

where $\tilde{G} = \sigma G$, and $\sigma \equiv \text{diag}\{1, -1\}$ is the diagonal matrix.

Following [26], let us assume that $G_0(x)$ is known which respects

$$i\frac{d}{dx}G_0(x) = G_0(x)M(x). \quad (10)$$

Being so, one is concerned, instead of (9), with solution of

$$\frac{d}{dx}(G_0(x)\tilde{G}(x, x')) = \frac{1}{i}e^{ix\xi}\delta(x - x') G_0(x). \quad (11)$$

Without going into details of [26], let us calculate

$$\mathcal{J}(x) = \iint e^{-ix\xi} G(x, x') dx' d\xi$$

(ξ -integration is also from $-\infty$ to ∞). Notice that $G(x, x')$ depends on ξ due to (8). Integrating (11) one obtains:

$$\begin{aligned} \mathcal{J}_{11}(x) = & \frac{1}{i}(\det G_0)^{-1} \int \left[G_0^{22}(x) \int_{+\infty}^x G_0^{11}(s) e^{i\xi(s-x)} ds \right. \\ & \left. - G_0^{12}(x) \int_{-\infty}^x G_0^{21}(s) e^{i\xi(s-x)} ds \right] d\xi, \end{aligned} \quad (12)$$

where the entries G_0^{ij} respect (10). Equations (11), (12) tell us that inhomogeneous Eq.(7) is solved, in fact, by the method of variation of arbitrary constant [29].

In its turn, (10) can be reduced to the couple of second order (Weber's, [3]) equations as follows. Let us note a useful matrix relation:

$$M^T - i\frac{d}{dx} = i\sigma u \begin{pmatrix} a^+ & \lambda \\ \lambda^* & -a^- \end{pmatrix} (\sigma u)^{-1}, \quad u \equiv \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \in U(2, \mathcal{C}), \quad (13)$$

where $\sigma \equiv \text{diag}\{1, -1\}$, $a^\pm \equiv x \mp d/dx$, $\lambda \equiv \omega + i\Delta$, and superscripts T and $*$ imply transposed and complex conjugated, respectively. Taking G_0 in the form

$$G_0(x) = \sqrt{2} \begin{pmatrix} h_1 & f_1 \\ h_2 & f_2 \end{pmatrix} u^{-1}, \quad (14)$$

and using

$$\begin{pmatrix} a^+ & \lambda \\ \lambda^* & -a^- \end{pmatrix} \begin{pmatrix} a^- & \lambda \\ \lambda^* & -a^+ \end{pmatrix} = \begin{pmatrix} a^+a^- + |\lambda|^2 & 0 \\ 0 & a^-a^+ + |\lambda|^2 \end{pmatrix},$$

one obtains unknown $f_{1,2} \equiv f_{1,2}(x)$ and $h_{1,2} \equiv h_{1,2}(x)$ from the couple of Weber's equations:

$$\frac{d^2}{dy^2}\tilde{f} - \left(\frac{y^2}{4} - \frac{1}{2} + |\lambda|^2\right)\tilde{f} = 0, \quad \frac{d^2}{dy^2}\tilde{h} - \left(\frac{y^2}{4} + \frac{1}{2} + |\lambda|^2\right)\tilde{h} = 0, \quad (14')$$

where $y \equiv x\sqrt{2}$, the tilde implies that f, h are expressed through y , and 1, 2 are omitted. Besides, λ is also rescaled in (14'), i.e., $\lambda \equiv (\omega + i\Delta)/\sqrt{2}$ until the end of the paper.

Eventually, the following combinations for $\tilde{h}_{1,2}$ and $\tilde{f}_{1,2}$ are chosen in [26] to express G_0^{ij} which ensure the integration in (12):

$$\tilde{h}_1 \pm \tilde{f}_1 = \mathcal{U}_+(y) \mp \lambda^{-1}\mathcal{U}_-(y), \quad \pm i(\tilde{f}_2 \mp \tilde{h}_2) = \mathcal{U}_+(-y) \mp \lambda^{-1}\mathcal{U}_-(-y),$$

where $\mathcal{U}_\pm(y) \equiv \mathcal{U}(|\lambda|^2 \pm 1/2, y)$ are the parabolic cylinder functions (the notation $\mathcal{U}(a, y)$ implies $D_{-a-\frac{1}{2}}(y)$). The asymptotical behaviour of $\mathcal{U}(a, y)$ is given by the estimates [3]:

$$\mathcal{U}(a, y) \simeq \exp(-y^2/4) y^{-a-1/2}, \quad \mathcal{U}(a, -y) \simeq \frac{(2\pi)^{1/2}}{\Gamma(a+1/2)} \exp(y^2/4) y^{a-1/2},$$

at $y \rightarrow +\infty$, $|y| \gg a$ (the leading terms).

The theory of distributions allows to calculate (12) straightforwardly. For the given choice of G_0^{ij} , the order of integrations in (12) can be changed at fixed x (Fubini's theorem). Integrating over ξ first, we obtain $2\pi\delta(s-x)$. Then, using the properties of δ -function [29] we obtain:

$$\mathcal{J}_{11}(x) = i\pi \frac{G_0^{11}(x) G_0^{22}(x) + G_0^{12}(x) G_0^{21}(x)}{G_0^{11}(x) G_0^{22}(x) - G_0^{12}(x) G_0^{21}(x)}$$

(again in the x -notations). A note about $\mathcal{J}_{21}$ will be given in the last section. Other entries are not of interest for us. The final answer for $\mathcal{J}_{11}(x)$ reads:

$$\begin{aligned} \mathcal{J}_{11}(x) = & \frac{1}{i} \left(\frac{\pi}{2}\right)^{1/2} \left(\lambda \Gamma(|\lambda|^2 + 1) \mathcal{U}_+(x\sqrt{2}) \mathcal{U}_+(-x\sqrt{2}) \right. \\ & \left. + \lambda^* \Gamma(|\lambda|^2) \mathcal{U}_-(x\sqrt{2}) \mathcal{U}_-(-x\sqrt{2}) \right), \end{aligned} \quad (15)$$

where $\Gamma(\cdot)$ is the gamma function [2].

3 THE SERIES REPRESENTATION

Let us consider another approach to (7), (8) [27]. The operator $\mathcal{H}(x)$ (7) is self-adjoint on $\mathcal{R}$, and it can be conjugated by the constant matrix u (13):

$$u^{-1} \mathcal{H} u = \mathcal{H}_{em}, \quad \mathcal{H}_{em} = \begin{pmatrix} \Delta & ia^- \\ -ia^+ & -\Delta \end{pmatrix}.$$

Squared operator $\mathcal{H}_{em}^2$ resembles the Hamiltonian of a spinning electron in a constant homogeneous magnetic field [28]. It is straightforward to obtain eigenvalues $E_0, \pm E_n$ and eigenfunctions $\hat{\Psi}_0, \hat{\Psi}_n^\pm$ ($n \geq 1$) of $\mathcal{H}_{em}(x)$:

$$\hat{\Psi}_0 = \begin{pmatrix} 0 \\ \psi_0(x) \end{pmatrix}, \quad E_0 = -\Delta,$$

$$\hat{\Psi}_n^{(s)} = \frac{1}{\sqrt{2E_n}} \begin{pmatrix} \sqrt{E_n + s\Delta} \psi_{n-1}(x) \\ -is\sqrt{E_n - s\Delta} \psi_n(x) \end{pmatrix}, \quad sE_n,$$

where $s = \pm$, $E_n = \sqrt{\Delta^2 + 2n}$, and $\psi_n(x)$ are the Hermite functions, $\psi_n(x) = \pi^{-1/4} \times (2^n n!)^{-1/2} e^{-x^2/2} H_n(x)$ [30]. Orthonormality and completeness of the system $\hat{\Psi}_0, \hat{\Psi}_n^\pm$ ($n \geq 1$) can be directly verified since $\psi_n(x)$ ($n \geq 0$) are orthogonal and complete in $L^2(\mathbb{R})$.

Let us take unknown G in the form (14) and pass to the equation

$$(i\omega + \mathcal{H}_{em}) \begin{pmatrix} h \\ f \end{pmatrix} = \delta(x - x') e^{ix\xi} \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$$

(the first column of the matrix equation in question). Now the unknown h and f depend on x and x' , and we expand $\begin{pmatrix} h \\ f \end{pmatrix}$ in $\hat{\Psi}_0(x), \hat{\Psi}_n^\pm(x)$ with the coefficients dependent on x' :

$$\begin{pmatrix} h \\ f \end{pmatrix}(x, x') = b(x') \hat{\Psi}_0(x) + \sum_{s=+,-} \sum_{n=1}^{\infty} b_n^{(s)}(x') \hat{\Psi}_n^{(s)}(x).$$

We define the coefficients [34] and, using

$$G_{11}(x) = \int dx' (h(x, x') + f(x, x')),$$

obtain $\mathcal{J}_{11}$ in the series form:

$$\mathcal{J}_{11}(x) = \frac{1}{i} \frac{\pi}{\sqrt{2}} \sum_{n=0}^{\infty} \psi_n^2(x) \left(\frac{\lambda}{|\lambda|^2 + n + 1} + \frac{\lambda^*}{|\lambda|^2 + n} \right). \quad (16)$$

In [26, 27] one is concerned with (15) and (16) themselves which correspond to the integrated entries of $G(x)$ (7). Let us now equate separately the real and imaginary parts of (15) and (16). After the analytical continuation $|\lambda|^2 \rightarrow z$, we deduce:

$$\Gamma(z) \mathcal{U}\left(z - \frac{1}{2}, x\sqrt{2}\right) \mathcal{U}\left(z - \frac{1}{2}, -x\sqrt{2}\right) = \sqrt{\pi} \sum_{n=0}^{\infty} \frac{\psi_n^2(x)}{n + z}, \quad \text{Re } z > 0. \quad (17)$$

The series in R.H.S. of (17) is convergent absolutely and uniformly on $\mathbb{R}$, and it is straightforward to express it in the integral form using

$$(n + z)^{-1} = \int_0^{\infty} e^{-t(n+z)} dt, \quad \text{Re } z > 0.$$

Using the generating function of Hermite polynomials [2] to sum up after exchanging summation and integration, one just obtains Eq.(6):

$$\begin{aligned} \Gamma(z) \mathcal{U}\left(z - \frac{1}{2}, x\right) \mathcal{U}\left(z - \frac{1}{2}, -x\right) &= \\ &= \frac{1}{\sqrt{2}} \int_0^{\infty} \exp\left(t\left(\frac{1}{2} - z\right) - \frac{x^2}{2} \tanh \frac{t}{2}\right) \frac{dt}{\sqrt{\sinh t}}, \quad \text{Re } z > 0, x \in \mathbb{R} \end{aligned} \quad (18.1)$$

(x is rescaled, $x\sqrt{2} \rightarrow x$). R.H.S. of (18.1) can be transformed to the integral which is known [6, 31]:

$$\int_0^1 \frac{(1-s)^{z-1}}{(1+s)^z} \exp\left(-\frac{x^2}{2}s\right) \frac{ds}{\sqrt{s}} = B\left(\frac{1}{2}, z\right) \Phi_1\left(\frac{1}{2}, z, z + \frac{1}{2}; -1, \frac{x^2}{2}\right), \quad \text{Re } z > 0, \quad (18.2)$$

where $\Phi_1(\alpha, \beta, \gamma; u, v)$ is the confluent hypergeometric series in two variables [1, 32].

4 THE SPECIAL CASES

Two ways of solving (7) give us, naturally, two representations for the same fundamental solution $G(x)$. Generally, it is a powerful technique – to equate equivalent representations of a given Green function [33]. We shall consider three particular examples in order to argue directly the interesting reductions of (18) to the known facts. Equation (18.1) depends on two parameters, $z \in \mathcal{C}$ and $x \in \mathbb{R}$, and their admissible values constitute the domain $\mathcal{D} = \{\text{Re } z > 0\} \times \mathbb{R}$. So, we shall consider the following three subsets in $\mathcal{D}$: the line, $z = 1/2$; open half-plane, $x = 0$; asymptotic regions, $z \gg 1$, $|x| \neq 0$ is bounded, and $|z|$ is limited, $|x| \gg 1$.

Example 1: $z = \frac{1}{2}, x \in \mathbb{R}$.

Using [2], we proceed:

$$\begin{aligned}\mathcal{U}(0, x) &\equiv D_{-\frac{1}{2}}(x) = \left(\frac{x}{2\pi}\right)^{1/2} K_{\frac{1}{4}}\left(\frac{x^2}{4}\right), \\ \mathcal{U}(0, -x) &\equiv D_{-\frac{1}{2}}(-x) = -i \left(\frac{x}{2\pi}\right)^{1/2} K_{\frac{1}{4}}\left(\frac{x^2}{4}\right) + i \left(\frac{x}{\pi}\right)^{1/2} K_{\frac{1}{4}}\left(-\frac{x^2}{4}\right).\end{aligned}$$

In their turn,

$$\begin{aligned}K_{\frac{1}{4}}\left(\frac{x^2}{4}\right) &= \frac{\pi}{\sqrt{2}} \left(I_{-\frac{1}{4}}\left(\frac{x^2}{4}\right) - I_{\frac{1}{4}}\left(\frac{x^2}{4}\right) \right), \\ K_{\frac{1}{4}}\left(-\frac{x^2}{4}\right) &= \frac{\pi}{2} \left((1-i) I_{-\frac{1}{4}}\left(\frac{x^2}{4}\right) - (1+i) I_{\frac{1}{4}}\left(\frac{x^2}{4}\right) \right).\end{aligned}$$

Then,

$$\mathcal{U}(0, -x) = \frac{(\pi x)^{1/2}}{2} \left(I_{-\frac{1}{4}}\left(\frac{x^2}{4}\right) + I_{\frac{1}{4}}\left(\frac{x^2}{4}\right) \right),$$

and

$$\pi^{1/2} \mathcal{U}(0, x) \mathcal{U}(0, -x) = \frac{\pi^{1/2} x}{2^{3/2}} K_{\frac{1}{4}}\left(\frac{x^2}{4}\right) \left(I_{\frac{1}{4}}\left(\frac{x^2}{4}\right) + I_{-\frac{1}{4}}\left(\frac{x^2}{4}\right) \right). \quad (19)$$

On the other hand, the L.H.S. of (18.2) can be written at $z = \frac{1}{2}$ as

$$\int_0^1 \exp\left(-\frac{x^2}{2}s\right) \frac{ds}{\sqrt{s(1-s^2)}}. \quad (20)$$

Accordingly to [31] (Chapter 2), the integral (20) is just equal to (19) (see APPENDIX).

Example 2: $\text{Re } z > 0, x = 0$.

At $x = 0$ the L.H.S. of (18.1) is [2]:

$$\Gamma(z) \mathcal{U}^2\left(z - \frac{1}{2}, 0\right) \equiv \Gamma(z) D_{-z}^2(0) = \frac{\sqrt{\pi}}{2} \frac{\Gamma\left(\frac{z}{2}\right)}{\Gamma\left(\frac{z+1}{2}\right)}.$$

R.H.S. of (18.1) takes at $x = 0$ the form of the definition of the beta function [2]:

$$\frac{1}{2} \int_0^1 s^{z/2-1} (1-s)^{-1/2} ds = \frac{1}{2} B\left(\frac{z}{2}, \frac{1}{2}\right), \quad \operatorname{Re}\left(\frac{z}{2}\right) > 0.$$

The coincidence is clear.

Example 3: $z \gg 1$, $|x| \neq 0$ is bounded, or $|z|$ is bounded, $|x| \gg 1$.

In leading order, we estimate the triple product $\Gamma(z)\mathcal{U}(z - \frac{1}{2}, x)\mathcal{U}(z - \frac{1}{2}, -x)$ in the L.H.S. of (18.1) as [3]:

$$\Gamma(z)\mathcal{U}\left(z - \frac{1}{2}, x\right)\mathcal{U}\left(z - \frac{1}{2}, -x\right) \simeq \left(\frac{2\pi}{x^2 + 4z - 2}\right)^{1/2}, \quad (21)$$

where $z > 1/2$, and $x^2 + 4z - 2$ is assumed positive and large. By steepest descent we estimate the R.H.S. of (18.1) at $z \gg 1$ ($|x| \neq 0$ is bounded) as $(\pi/(2z))^{1/2}$. Analogously, the integral in the L.H.S. of (18.2) can be estimated at $|x| \gg 1$ ($|z|$ is bounded) as $(2\pi)^{1/2}/|x|$. In the both cases we get agreement with (21). However, when z and $|x|$ are large, (18.1) results in (21) as well.

5 CONCLUSION

To conclude, the following remarks are in order. The relation

$$D_\nu(z) = 2^{\frac{\nu}{2} + \frac{1}{4}} \frac{1}{\sqrt{z}} W_{\frac{\nu}{2} + \frac{1}{4}, \pm \frac{1}{4}}\left(\frac{z^2}{2}\right)$$

is known which relates $D_\nu(z)$ to the Whittaker function. Thus, our product $D_\nu(z)D_\nu(-z)$ can be expressed by means of the integral representation for $W_{\kappa,\mu}(z)W_{\chi,\mu}(z')$ [2] (though containing the hypergeometric function ${}_2F_1$ under integration). However, some extra work is still required to relate (6) to the corresponding formula. Consideration of the entry $\mathcal{J}_{21}$ does not give really different formula: we obtain a representation which can be deduced if one applies $x - d/dx$ to (6) (compare with (3)). The representation (6) looks simpler in comparison with analogous formulas mentioned above, and hopefully can be useful in applications.

ACKNOWLEDGEMENTS

The author is grateful to Prof. V. M. Babich for useful discussions. The research described has been supported in part by RFBR, No 01-01-01045

APPENDIX

More precisely, [31] provides the formula for the integral

$$\int_0^a \frac{x^{n-\frac{1}{2}}}{\sqrt{a^2 - x^2}} e^{-px} dx,$$

which can be obtained from

$$\begin{aligned} \int_0^a \frac{e^{-px} dx}{\sqrt{x(a^2 - x^2)}} &= \frac{(\pi p)^{1/2}}{2} K_{\frac{1}{4}}\left(\frac{ap}{2}\right) \\ &\times \left(I_{\frac{1}{4}}\left(\frac{ap}{2}\right) + I_{-\frac{1}{4}}\left(\frac{ap}{2}\right) \right), \quad a > 0, \operatorname{Re} p > 0 \end{aligned} \quad (a)$$

by multiple differentiation $\partial^n / \partial(-p)^n$. Besides, there exists another integral formula [31]:

$$\begin{aligned} \int_a^\infty \frac{e^{-px} dx}{\sqrt{x(x^2 - a^2)}} &= \frac{(\pi p)^{1/2}}{2} K_{\frac{1}{4}}\left(\frac{ap}{2}\right) \\ &\times \left(I_{-\frac{1}{4}}\left(\frac{ap}{2}\right) - I_{\frac{1}{4}}\left(\frac{ap}{2}\right) \right), \quad a > 0, \operatorname{Re} p > 0. \end{aligned} \quad (b)$$

The couple of Eqs.(a) and (b) can be related to the following couple of the integral transforms [35]:

$$\begin{aligned} \int_0^\infty \frac{\sin bx dx}{\sqrt{x(x^2 + z^2)}} &= \left(\frac{\pi b}{2}\right)^{1/2} K_{\frac{1}{4}}\left(\frac{bz}{2}\right) I_{\frac{1}{4}}\left(\frac{bz}{2}\right), \quad b > 0, \operatorname{Re} z > 0, \\ \int_0^\infty \frac{\cos bx dx}{\sqrt{x(x^2 + z^2)}} &= \left(\frac{\pi b}{2}\right)^{1/2} K_{\frac{1}{4}}\left(\frac{bz}{2}\right) I_{-\frac{1}{4}}\left(\frac{bz}{2}\right), \quad b > 0, \operatorname{Re} z > 0. \end{aligned}$$

Notes added after release of the paper

1.

It is appropriate to point out that the representation

$$\begin{aligned} \exp(-x^2) \left(H_\lambda^2(x) + G_\lambda^2(x) \right) &= 2^{\lambda+1} \pi^{-1} \Gamma(\lambda + 1) \\ &\times \int_0^\infty \exp\left(-(2\lambda + 1)t + x^2 \tanh t\right) (\cosh t \sinh t)^{-1/2} dt \end{aligned} \quad (N1)$$

has been found in [36]. Here, H_λ and G_λ are the Hermite functions which are related to the confluent hypergeometric function $\Phi(a, b; z)$,

$$\Phi(a, b; z) = \frac{\Gamma(b)}{\Gamma(a)} \sum_{n=0}^\infty \frac{\Gamma(a+n)}{\Gamma(b+n)} \frac{z^n}{n!},$$

as follows [37]:

$$\begin{aligned} 2^{-\lambda} H_\lambda(x) &= \frac{\Gamma(1/2)}{\Gamma(1/2 - \lambda/2)} \Phi\left(-\frac{\lambda}{2}, \frac{1}{2}; x^2\right) + \frac{\Gamma(-1/2)}{\Gamma(-\lambda/2)} x \Phi\left(-\frac{\lambda}{2} + \frac{1}{2}, \frac{3}{2}; x^2\right), \\ 2^{-\lambda} G_\lambda(x) &= -\frac{\Gamma(1/2)}{\Gamma(1/2 - \lambda/2)} \tan \frac{\pi\lambda}{2} \Phi\left(-\frac{\lambda}{2}, \frac{1}{2}; x^2\right) + \frac{\Gamma(-1/2)}{\Gamma(-\lambda/2)} x \cot \frac{\pi\lambda}{2} \Phi\left(-\frac{\lambda}{2} + \frac{1}{2}, \frac{3}{2}; x^2\right). \end{aligned}$$

Equation (N1) appears in [36] as a confluent limit of an integral representation for product of two Gegenbauer functions. The representation (N1) turns out to be useful [36] in the problem of monotonicity of differences of zeros of the Hermite functions [38, 39].

Using connection [37] between $\Phi(a, b; z)$ and the parabolic cylinder functions, one finds:

$$\exp(-x^2/2) H_\lambda(x) = 2^{\lambda/2} D_\lambda(x\sqrt{2}), \quad (N2)$$

$$\exp(-x^2/2) G_\lambda(x) = \frac{2^{\lambda/2}}{\sin \pi \lambda} \left(\cos \pi \lambda D_\lambda(x\sqrt{2}) - D_\lambda(-x\sqrt{2}) \right).$$

Eventually, using (N2) and the linear relations [2] between the parabolic cylinder functions, it is straightforward to check that (N1) and (6) (and thus (18)) are equivalent.

2.

Equation analogous to (17) can also be written for the product of two parabolic cylinder functions at coinciding arguments:

$$\Gamma(z) \mathcal{U}^2 \left(z - \frac{1}{2}, x\sqrt{2} \right) = \sqrt{\pi} \sum_{n=0}^{\infty} \frac{(-1)^n \psi_n^2(x)}{n+z}, \quad \text{Re } z > 0, \quad (N3)$$

where summation in R.H.S. goes over simple poles of $\Gamma(z)$. In the same way as in Sec.3, we obtain from (N3):

$$D_{-z}^2(x) = 2^{-1/2} \Gamma^{-1}(z) \int_0^\infty \exp \left(\left(\frac{1}{2} - z \right) t - \frac{x^2}{2} \coth \frac{t}{2} \right) \frac{dt}{\sqrt{\sinh t}}, \quad \text{Re } z > 0, \quad (N4)$$

or

$$D_{-z}^2(x) = \Gamma^{-1}(z) \int_1^\infty \frac{(s-1)^{z-1}}{(s+1)^z} \exp \left(-\frac{x^2}{2} s \right) \frac{ds}{\sqrt{s}}. \quad (N5)$$

However, the representation (N5) is known [31] just for the square of the parabolic cylinder function but in another form: i.e., for the integration variable obtained by shift $s-1 \rightarrow s$. The integral in (18.2) and Eq.(N5) imply the following combinations:

$$D_{-z}(x) (D_{-z}(x) + D_{-z}(-x)) = \Gamma^{-1}(z) \int_0^\infty \frac{|s-1|^{z-1}}{(s+1)^z} \exp \left(-\frac{x^2}{2} s \right) \frac{ds}{\sqrt{s}}, \quad (N6)$$

$$D_{-z}(x) (D_{-z}(x) - D_{-z}(-x)) = \Gamma^{-1}(z) \int_0^\infty \frac{|s-1|^z}{(s+1)^z} \frac{1}{s-1} \exp \left(-\frac{x^2}{2} s \right) \frac{ds}{\sqrt{s}}.$$

Besides, (18.1) and (N4) can be re-written together in the following form:

$$D_{-z}(x) D_{-z}(\pm x) = \frac{\exp(-x^2/2)}{2^{1/2} \Gamma(z)} \int_0^\infty \exp \left(\left(\frac{1}{2} - z \right) t \mp x^2 (e^t \mp 1)^{-1} \right) \frac{dt}{\sqrt{\sinh t}}. \quad (N7)$$

The Author is grateful to Prof. M. E. Muldoon for attracting his attention to [36] and for interesting correspondence which inspired to write down Eqs.(N3)–(N7) and to present the given notes.

References

- [1] H. Bateman, A. Erdélyi, *Higher Transcendental Functions*, (McGraw–Hill, New York, 1953/55).
- [2] W. Magnus, F. Oberhettinger, P. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, (Springer-Verlag, Berlin, 1966).
- [3] M. Abramowitz, I. A. Stegun (eds.), *Handbook of Mathematical Functions*, (Dover, New York, 1970).
- [4] J. C. P. Miller, *Tables of Weber Parabolic Cylinder Functions*. In: *Tables of Weber Functions*, K. A. Karpov, Ed. (Comp. Center Acad. Sci. USSR, Moscow, 1968) pp.9–87 [In Russian]
- [5] M. K. Kerimov, *Some New Results on the Theory of Weber Functions*. In: *Tables of Weber Functions*, K. A. Karpov, Ed. (Comp. Center Acad. Sci. USSR, Moscow, 1968) pp.88–116 [In Russian]
- [6] I. S. Gradstein, I. M. Ryzhik, *Tables of Integrals, Sums, Series, and Products*, (Nauka, Moscow, 1971) [In Russian]
- [7] C. S. Meijer, *Über die Integraldarstellungen der Whittakerschen Funktion $W_{k,m}(z)$ und der Hankelschen und Besselschen Funktionen*, Nieuw Archief voor Wiskunde, **18**, No 2 (1934), 35–57.
- [8] C. S. Meijer, *Noch einige Integraldarstellungen für die Whittakersche Funktion*, Proc. Konink. Akad. Wetens. Amst., **38**, No 5 (1935), 528–535.
- [9] C. S. Meijer, *Über eine Integraldarstellung der Whittakerschen Funktion*, Proc. Konink. Nederl. Akad. Wetens. Amst., **41**, No 1 (1938), 42–44.
- [10] C. S. Meijer, *Einige Integraldarstellungen für Whittakersche und Besselsche Funktionen*, Proc. Konink. Akad. Wetens. Amst., **37**, No 10 (1934), 805–812.
- [11] C. S. Meijer, *Einige Integraldarstellungen für Produkte von Whittakersche Funktionen*, Quart. J. Math. (Oxford Ser.), **6**, No 24 (1935), 241–248.
- [12] C. S. Meijer, *Einige Integraldarstellungen aus der Theorie der Besselschen und Whittakerschen Funktionen*, Proc. Konink. Akad. Wetens. Amst., **39**, No 3 (1936), 394–403; No 4 (1936), 519–527.
- [13] C. S. Meijer, *Über Whittakersche bzw. Besselsche Funktionen und deren Produkte*, Nieuw Archief voor Wiskunde, **18**, No 4 (1936), 10–39.
- [14] C. S. Meijer, *Über Produkte von Whittakerschen Funktionen*, Proc. Konink. Akad. Wetens. Amst., **40**, No 2 (1937), 133–141; No 3 (1937), 259–263.
- [15] C. S. Meijer, *Note über das Produkt $M_{k,m}(z) M_{-k,m}(z)$* , Proc. Konink. Nederl. Akad. Wetens. Amst., **41**, No 3 (1938), 275–277.

- [16] W. N. Baily *An Integral Representation for the Product of Two Whittaker Functions*, Quart. J. Math. (Oxford Ser.), **8**, No 29 (1937), 51–53.
- [17] S. C. Dhar, *On the Product of Parabolic Cylinder Functions*, J. Indian Math. Soc. (N.S.), **1**, No 3 (1934), 105–108.
- [18] S. C. Dhar, *On the Product of Parabolic Cylinder Functions with Different Arguments*, J. London Math. Soc., **10**, No 39 (1935), 171–175.
- [19] V. L. Mutatker, *On Some Expansions and Integrals Involving the Parabolic Cylinder Functions*, J. Indian Math. Soc. (N.S.), **1**, No 2 (1934), 53–58.
- [20] S. C. Mitra, *On the Squares of Weber’s Parabolic Cylinder Functions and Certain Integrals Connected with Them*, Proc. Edinburgh Math. Soc. (Ser.2), **4**, Part I (1934) 27–32.
- [21] W. T. Howell, *On Some Operational Representations of Products of Parabolic Cylinder Functions and Products of Laguerre Polynomials*, Phil. Mag.(7), **24**, No 165 (1937), 1082–1093.
- [22] W. T. Howell, *Integral Representations for Products of Weber’s Parabolic Cylinder Functions*, Phil. Mag.(7), **25**, No 168 (1938), 456–458.
- [23] H. Shanker, *On Certain Integrals and Expansions Involving Weber’s Parabolic Cylinder Functions*, J. Indian Math. Soc. (N.S.), **4**, No 4 (1940), 158–166.
- [24] E. Feldheim, *Expansions and Integral-Transforms for Products of Laguerre and Hermite Polynomials*, Quart. J. Math. (Oxford Ser.), **11**, No 41 (1940), 18–29.
- [25] V. M. Babich, D. B. Dement’ev, B. A. Samokish, V. P. Smyshlyaev, *Scattering of a Highfrequency Wave by the Vertex of an Arbitrary Cone. (Singular Directions.)*, In: Zap. Nauch. Semin. POMI, **264**, (2000), 7–21 [Transl.: J. Math. Sci.].
- [26] R. Combescot and T. Dombre, *Superfluid Current in $^3\text{He-A}$ at $T = 0$* , Phys. Rev. B, **28**, No 9 (1983), 5140–5148.
- [27] C. Malyshev, *Higher Corrections to the Mass Current in Weakly Inhomogeneous Superfluid $^3\text{He-A}$* , Phys. Rev. B, **59**, No 10 (1999), 7064–7075.
- [28] L. D. Landau, E. M. Lifchitz, *Quantum Mechanics*, (Nauka, Moscow, 1989) [In Russian]
- [29] V. S. Vladimirov, *Equations of Mathematical Physics*, (Nauka, Moscow, 1971) [In Russian]
- [30] A. N. Tikhonov, A. A. Samarskij, *Equations of Mathematical Physics*, (Nauka, Moscow, 1966) [Transl.: New York, Pergamon, 1963]
- [31] A. P. Prudnikov, Ju. A. Brychkov, O. I. Marichev, *Integrals and Series. Elementary Functions*, (Nauka, Moscow, 1981) [Transl.: New York, Gordon and Breach]
- [32] A. P. Prudnikov, Ju. A. Brychkov, O. I. Marichev, *Integrals and Series. Additional Chapters*, (Nauka, Moscow, 1986) [In Russian]

- [33] P. M. Morse, H. Feshbach, *Methods of Theoretical Physics*, (McGraw-Hill, New York, 1953)
- [34] C. Malyshev, *On the Two Calculations of the Superfluid Current in the A-Phase of Helium-3*, In: Zap. Nauch. Semin. POMI, **209**, (1994), 179–193 [Transl.: J. Math. Sci., **83**, (1997), 113].
- [35] H. Bateman, A. Erdélyi, *Tables of Integral Transforms*, (McGraw-Hill, New York, 1954).
- [36] L. Durand, *Nicholson-Type Integrals for Products of Gegenbauer Functions and Related Topics*, In: Theory and Applications of Special Functions, Richard A. Askey, Ed. (Academic Press, New York, etc., 1975) pp.353–374
- [37] D. S. Kuznetsov, *Special Functions*, (Highest School, Moscow, 1965) [In Russian]
- [38] L. Lorch, P. Szego *Higher Monotonicity Properties of Certain Sturm–Liouville Functions*, Acta Math., **109**, No 1-2 (1963), 55–73.
- [39] L. Lorch, M. E. Muldoon, P. Szego *Higher Monotonicity Properties of Certain Sturm–Liouville Functions III, IV*, Can. J. Math., **22**, No 6 (1970), 1238–1265; **24**, No 2 (1972), 349–368.